200500102

## HHE UNITED SHATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

South Ankota Agricultural Experiment Station

THETERS, THERE HAS BEEN PRESENTED TO THE

#### Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT. THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE THEREOF IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED BY THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE PLENT TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFFERING IT FOR SALE, OR REPRODUCING IT, OR CONTING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE SEED OF THE ABOVE BE OR USING IT IN PRODUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT OF SAID BY VARIETY NAME ONLY AS A CLASS OF CERTIFIED SEED AND (2) SHALL CONFORM TO THE NUMBER OF THE RIGHTS. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321

WHEAT, COMMON

'Wendy'

In Jestimon Macrest, I have hereunto set my hand and caused the seal of the Plant Anciety Frotection Office to be affixed at the City of Washington, D.C. this sixth day of Warch, in the year two thousand and six.

Attest:

Commissioner

Plant Variety Protection Office Agricultural Marketing Service Agriculture

The following statements are made in accordance with the Privacy Act of 1974 (5 U.S.C. 552a) and the Paperwork Reduction Act (PRA) of 1995.

Application is required in order to determine if a plant variety protection certificate is to be issued (7 U.S.C. 2421). Information is held confidential until certificate is issued (7 U.S.C. 2426).

### U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE AND TECHNOLOGY - PLANT VARIETY PROTECTION OFFICE APPLICATION FOR PLANT VARIETY PROTECTION CERTIFICATE (Instructions and information collection burden statement on reverse) 1. NAME OF OWNER 2. TEMPORARY DESIGNATION OR 3. VARIETY NAME

	•			EXPERIMENTAL NAME							
South Dakota Agri	cultural Exper	iment	Station	SD97W604	Wendy						
4. ADDRESS (Street and No., or R.F.D. No.	., City, State, and ZIP Code, and Count	try)		5. TELEPHONE (include area code)	FOR OFFICIAL USE ONLY						
South Dakota Stat	e University			(605) 688-4149	新華新 Dat 特殊議員を記すると 日 八丁						
Ag Hall 129					PVPO NUMBER						
Brookings SD 57	007			6. FAX (include area code)	00500102						
•				(605) 688-6065	FILING DATE						
7. IF THE OWNER NAMED IS NOT A "PERS	SON", GIVE FORM OF	8. IF INCOR	PORATED, GIVE	9. DATE OF INCORPORATION	,						
ORGANIZATION (corporation, partnership	o, association, etc.)	STATE OF	INCORPORATION								
Agricultural Expe	riment Station	1.	N/A	N/A	3005,18.NAC						
10. NAME AND ADDRESS OF OWNER REP	PRESENTATIVE(S) TO SERVE IN THIS	S APPLICATION	l. (First person listed will re	eceive all papers)	FILING AND EXAMINATION FEES:						
Dr. Kevin Kephart				m, W. Wheat Breed	2E						
SD Ag Exp Station South Dakota State			Science D		\$ 3652.00						
Brookings SD 5700	<u></u>			140c, SDSU	E DATE 1/3/105						
	• •	DIOOK	rugs su 5	7007	1131/02						
		· ·		,	V CERTIFICATION FEE:						
					. 768.00						
					DATE 2/14/06						
11. TELEPHONE (Include area code)	12. FAX (Include area code)	13.	E-MAIL	14. CROP	KIND (Common Name)						
(605) 688-4453	605-688-4452	An	ir.Ibrahin	n@sdstate.edu Har	d white winter						
			· I'		wheat						
<ol> <li>CHECK APPROPRIATE BOX FOR EACH reverse)</li> </ol>	ATTACHMENT SUBMITTED (Follow	instructions on	19. DOES THE O	WNER SPECIFY THAT SEED OF THIS VAI SEED? See Section 83(a) of the Plant Va							
a. 🗵 Exhibit A. Origin and Breeding	History of the Variety		ICK YES	6 (If 'yes', answer items 20	NO (if "no," go to item 22)						
b. X Exhibit B. Statement of Distinct	tness			and 21 below)							
c. Exhibit C. Objective Description	n of Variety			WNER SPECIFY THAT SEED OF THIS LIMITED AS TO NUMBER OF CLASSES?	$\square$ YES $XX^\square$ NO						
d. 🗵 Exhibit D. Additional Descriptio	n of the Variety (Optional)		IF YES, WHICH CLASSES?    FOUNDATION    REGISTERED    CERTIFIED								
e. 😾 Exhibit E. Statement of the Bas											
f. X Voucher Sample (2,500 viable to verification that tissue culture we repository)	untreated seeds or, for tuber propagate ill be depositied and maintained in an a	ad varieties, approved public		WNER SPECIFY THAT THE CLASSES BE TO NUMBER OF GENERATIONS?	YES NO						
<u>x</u> .	.705), made payable to "Treasurer of th	he United	IF YES, SPEC								
States" (Mail to the Plant Variet	,705), made psyable to "Treasurer of the y Protection Office)		NUMBER 1, 2	, 3, etc. FOUNDATION	REGISTERED CERTIFIED						
			(If additional e	xplanation is necessary, please use the spa	ce indicated on the reverse.)						
2. HAS THE VARIETY (INCLUDING ANY HA	RVESTED MATERIAL) OR A HYBRID	PRODUCED	23. IS THE VARIE	TY OR ANY COMPONENT OF THE VARIE	TY PROTECTED BY INTELLECTUAL						
FROM THIS VARIETY BEEN SOLD, DISP OTHER COUNTRIES?	OSED OF, TRANSFERRED, OR USE	D IN THE U. S.		RIGHT ( <i>PLANT BREEDER'S RIGHT OR PA</i> 1	•						
☐ YES	XX NO		'		•						
IF YES, YOU MUST PROVIDE THE DATE	OF FIRST SALE, DISPOSITION, TR	ANSFER, OR U	SE REFERENCE I	COUNTRY, DATE OF FILING OR ISSUAND NUMBER. <i>(Please use space indicated on I</i>							
FOR EACH COUNTRY AND THE CIRCU	MSTANCES. (Please use space indica	ated on reverse.	<u> </u>								
<ol> <li>The owners declare that a viable sample of for a tuber propagated variety a tissue cult</li> </ol>	f basic seed of the variety will be furnis ure will be deposited in a public reposi	thed with applications and mainta	ation and will be replenished	ed upon request in accordance with such reg certificate.	rulations as may be applicable, or						
				at the variety is new, distinct, uniform, and s							
Owner(s) is(are) informed that faise repres					•						
			portation.								
IGNATURE OF OWNER			SIGNATURE OF	OWNER							
AME (Please print or type)			11112								
	in D. Kephart		NAME (Please pri	nt or type)	•						
DI. KEV.	rn p. Kehnart			·							
APACITYORIME irector. SD Ag Ext	OATE	1/2005	CAPACITY OR TI	TLE	DATE						

## EXHIBIT A Wendy (SD97W604) Origin and Breeding History of the Variety

Wendy was derived from the cross SD89333/'Abilene' made in 1992. SD89333 is an unreleased experimental line from South Dakota State University with the pedigree 'Gent'/'Siouxland'. Wendy was developed by means of the bulk breeding method. The cross (coded X92259) was advanced to the F<sub>3</sub> generation as a bulk. Seed harvested from the F<sub>3</sub> bulk was sorted for white kernel color in 1995. The bulk of selected white kernels was coded X92259W and was grown in the greenhouse in 1996. Single heads were harvested from the selected F<sub>4</sub> bulk and planted in the field as head-rows in fall 1996. Wendy was derived as an F<sub>4.5</sub> line selected by S.D. Haley in 1997. Wendy was evaluated as SD97W604 in the South Dakota Early Yield Trial nursery in 1998. It was advanced beyond the Preliminary Yield Trial to the South Dakota Advanced Yield Trial in 1999 due to superior performance. It was tested in the South Dakota Crop Performance Testing (CPT) Variety Trial between 2000 and 2004, in the Northern Regional Performance Nursery during 2001 and 2002, and in the Southern Regional Performance Nursery in 2004.

Composite milling and bread baking properties of Wendy were determined in 2001 and 2002 in cooperative baking tests conducted by the USDA/ARS Hard Winter Wheat Quality Laboratory in Manhattan, KS.

Breeder seed of Wendy originated from a composite of 200 F<sub>10:11</sub> head-rows selected in 2002 based on visual uniformity and white kernel color purity. Wendy has been uniform for all morphological characters (such as maturity and plant height) during the last four generations of selfing and increase. Wendy contains 0.14% hard red grain. It also contains red and tall white off-types in the frequency of 0.05% and 0.01%, respectively.

Pen Phone conversation 1/24/2006

## EXHIBIT B Wendy (SD97W604) Statement of Distinctness


- 'Wendy' is most similar to the hard white winter wheat cultivar 'Trego', but differs in the following characteristics:
- 1) Grain Yield: Wendy has 0.7 LB/ACRE more grain yield than Trego, when recorded directly from the combine (Table 1).
- 2) Test Weight: Wendy has 0.3 LB/Bu less test weight than Trego, when recorded directly from the combine (Table 1).
- 3) Heading Date: Wendy heading is earlier by two days than Trego (Table 1).
- 4) Plant Height: Wendy is 2.0 inches shorter than Trego (Table 1).
- 5) Acid Polyacrylamide Gel Electrophoresis (PAGE): Acid Polyacrylamide Gel Electrophoresis shows that Wendy, Trego and 'Nuplains' have distinctly different seed storage protein banding patterns (Photograph 1). The arrows on the photograph point to bands present in Expedition but missing in Alliance and vice versa.
- 6) Urea Polyacrylamide Gel Electrophoresis: Urea Polyacrylamide Gel Electrophoresis also shows that Wendy, Trego and Nuplains have distinctly different seed storage protein banding patterns (Photograph 2). The arrows on the photograph point to bands present in Expedition but missing in Alliance and vice versa

Davis, B. 1964. Disc electrophoresis. II. Method and application to human serum protein. Ann. New York Acad. Sci. 121:404-427.


Table 1. South Dakota State University, Winter Wheat Breeding Trials Combined Over Locations and Years (2001 – 2003).

ÍĎ.	Grain yield (bu/a)	Test weight (lb/bu)	Heading (days)	Height (inches)
ARAPAHOE	50.1	58.2	159	33.0
CRIMSON	46.7	59.4	161	35.1
EXPEDITION	48.9	58.3	156	30.9
HARDING	48.3	58.5	161	35.2
NUPLAINS	46.3	59.2	161	29.6
WENDY	48.7	58.7	156	27.5
TREGO	48.0	59.0	158	29.5
Mean	48.7	58.4	159	31.8
CV%	13.2	3.2	0.7	4.8
LSD 0.05	1.6	0.5	0.5	0.6

Photograph I. Acid Polyacrylamide Gel Electrophoresis (PAGE) of hard white winter wheat cultivars Wendy, Trego, and Nuplains. The Acid PAGE was conducted by Dr. Brent Turnipseed, Seed Testing Lab, Plant Science Department, South Dakota State University.



Photograph 2. Urea Polyacrylamide Gel Electrophoresis (PAGE) of hard white winter wheat cultivars Wendy, Trego, and Nuplains. The Urea PAGE was conducted by Ms. Mary Thompson, Seed Testing Lab, Plant Science Department, South Dakota State University.



Plant Science Department

Box 2140C, NPB 247 SDSU Brookings, SD 57007-2141 Phone: 605-688-4020 FAX: 605-688-4452

#### PROCEDURE FOR POLYACRYLAMIDE GEL ELECTROPHORESIS

Written by J. Schneider and T. Machacek 1988 Revised 1989 by J. Johnson Revised 1991 by L. Fleege

This paper is written with the intent of passing on some practical information to laboratories which are interested in using polyacrylamide gel electrophoresis (PAGE) for the analysis of varietal purity of seed lots. Perhaps the practical experience we have gained from organizing and implementing our system will be of some assistance.

The lists which have been compiled are quite comprehensive of the needs for setting up a laboratory for PAGE. Personal preference may dictate changes in the glass and hardware, however the items listed seem to work the best for us. The equipment listed is essential for producing clear and reproducible results.

The chemical sources listed are those recommended to us (Lookhart et al., 1981) as the ones that produce the best results. Aluminum Lactate quality is important; it should be pure white (powdered sugar color). Impure Aluminum Lactate will result in blurred electrophorograms.

Reference samples of foundation seed (one pound) of each variety to be tested were collected and stored in plastic containers.

Thirty gram samples are collected in Udy bottles following purity testing. 100 seeds counted from each sample are ground in a small coffee grinder (Miracle Mill) and sifted through a 1.016 sieve. Use caution to avoid contamination between samples. Cleaning the grinder and sieve with forced air between samples is recommended.

#### PROTEIN EXTRACTION

Using a fine point marking pen, mark a 1.5 ml centrifuge tube for each sample. Wheat into each tube, add 750  $\mu$ l of Ethylene glycol and 0.25 g of flour. Extractions mix easier if the ethylene glycol is put in the tube first. Vortex the samples and wait an hour before centrifuging. Oat; into each tube, add 700  $\mu$ l of ethylene glycol and 0.30 g of flour. Vortex the samples and wait one hour before centrifuging. Single Seed: Crush each seed between weighing paper with a pliers. Place each into centrifuge tubes with 100  $\mu$ l of ethylene glycol and allow to set 1 hour. Load the centrifuge evenly. Run it on low speed (4500 x g recommended) for 10 minutes. [Optional: Add one drop of 10% methyl green dye (1 g methyl green brought to a volume of 10ml with  $H_1O$ ) to each sample with a small sized auto pipette].

#### MIXING THE GEL SOLUTION

Each 1.5 mm thick gel will require 50 ml of gel solution. Use a 125 ml vacuum flask with stopper when mixing 100 ml of gel solution. Into the flask add each of the following: Note: when working with the gel solution wear gloves and a mask as acrylamide is a neurotoxin until it has solidified.

For two gels: Aluminum Lactate (0.25g); Ascorbic Acid (0.024g); FeSO<sub>4</sub>-7H<sub>2</sub>O (200 $\mu$  of 10mg/10ml mixed fresh);

Wheat:

Bis-Acrylamide (0.25g), Acrylamide (6.00g);

<u> Oat:</u>

Bis-Acrylamide (0.4g), Acrylamide (7.50g). Add distilled water to obtain a

volume of 100ml.

Add a stir bar, stopper the flask, and place on the stirrer at slow speed, connect the vacuum and allow it to stir for 20 minutes. (The vacuum step is optional.) After this time is up, allow the solution to continue stirring while the pH is checked and adjusted. Lower the pH to 3.1 by titrating with Lactic Acid. If the pH is over adjusted, Trizma base (5M) can be used to increase the pH without adverse effects. Divide the solution into 2 portions in 100ml flasks containing stir bars.

Prepare the gel places making sure that they are tightly sealed to the rubber gaskets.

(This is very important but easy to forget).

Place one of the flasks containing gel solution on the stir plate at very slow speed. Note: excessive agitation mixes in oxygen which ties up the catalyst and prevents complete polymerization. Add 25  $\mu$ l of  $H_1O_1$  and allow to stir for 45 seconds. Carefully pour the solution between the plates until it is 1/4" from the top. Insert the gel comb making sure there are no bubbles trapped on the well ends. Repeat the procedure for the other gel and allow both to polymerize for 15 minutes.

#### TANK BUFFER

While waiting for the gel to polymerize, mix the tank buffer. Place 5.625 g of Aluminum Lactate in a 4000 ml flask. Fill the flask to approximately 4500 ml (middle of the neck on a 4000 ml flask) with distilled  $H_2O$  and stir. While the buffer stirs, check the pH. Titrate to 3.1 using Lactic Acid.

#### SAMPLE PLACEMENT

By this time the gel should be polymerized. Carefully remove the gel comb. Careless removal can cause suction, pulling the gel walls together and possibly breaking them. Pipette out any excess fluid in the slots using a long needle syringe. If the walls of the gel are crooked, they can be straightened at this time with the needle. Pill the slots about 2/3 full with buffer solution. Then, using a digital pipette, place 7.5  $\mu$ l of the sample extract ( $12\mu$ l for single seeds) into the slots. The extract should sink to the bottom. Using a pasteur pipette finish filling the slots with buffer.

RUNNING THE GEL

Attach the upper busser chamber to the gel plates making sure of a tight scal. Fill the upper busser chamber with busser one inch from the top. Pour the remaining busser into the lower busser chamber along with the stir bar. Set the tank on a stir plate and set the stir plate to a sast speed. This aids in maintaining an even temperature during the run.

Put the apparatus together. Make sure that the cathode is connected to the lower buffer and the anode to the top. Start the water bath and set it for 15C. Set the power source to constant current, switch it on and turn power up to 500 volts. The amperage, when running two gels, should initially read approximately 160 to 170 ma. Allow it to run 2

hours for wheat and oats.

#### MIXING THE STAINING SOLUTION

Take safety precautions to prevent eye and skin contact when working with the acid solutions. To make 600 ml of stain, mix in a 1000 ml flask the following: 100 grams TCA and 600 ml distilled  $H_2O$ . Add 20 ml of 1% Comassic brilliant blue (CBB) solution (1g CBB per 100ml ETOH).

#### REMOVING THE GELS

After 2 hours turn the power off and remove the gel plates. Remove the spacers and pry the plates apart (It is recommended using one of the spacers for this task. Take care as to not damage the spacer in the process). To keep track of the slot order, cut off the lower left hand corner. Then remove the slot fringes. The gel can then be loosened by running a spatula between it and the glass plate while squirting water in the gap created by the spatula. The gel should slide off into the staining pan (be sure to label the pan).

#### STAINING

Remove any excess water and add 300ml of stain. Place the trays on a shaker at gentle speed and allow them to stain for at least 4-6 hours. Over staining will not harm the gels or their results. Sometimes overnight staining is more convenient.

#### DESTAINING

Remove the stain (A handy way to remove liquids is with the vacuum pump. Put a catch bottle in the line and use a pasteur pipette in the hose end.) Add 300ml of  $H_2O$  to the tray. Put the trays on the shaker and allow to destain for 2-6 hours or whatever is necessary for acceptable results. A useful tool for moving the gel is a piece of plastic 11" x 6" from a report cover. Use the plastic sheet to place the gels into one quart zip-lock freezer bags.

#### CLEANING THE GEL PLATES

For easy and flawless removal of the gels, a thorough cleaning of the plates is essential. Wash them well with soapy water. Then wipe them down with ammonia glass cleaner or ethanol. Put the glass together with the spacers and clamps making sure that everything is flush on the bottom. Put them in the stand, insert the combs, and cover with plastic wrap. They are now ready for the next use.

REPRODUCE LOCALLY. Include form number and date on all reproductions.

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this collection of information is (0581-0055). The time required to complete this information collection is estimated to average 30 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact the USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call (202) 720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.

> U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE AND TECHNOLOGY PLANT VARIETY PROTECTION OFFICE BELTSVILLE, MD 20705

EXHIBIT C (Wheat)

#### **OBJECTIVE DESCRIPTION OF VARIETY** WHEAT (Triticum son )

NAME OF APPLICATION:  ADDRESS (Poord and No. or RD No., Cop., Suns, and Exp. Col.)  ADDRESS (Poord and No. or RD No., Cop., Suns, and Exp. Col.)  PLEASE READ ALL INSTRUCTIONS CAREFULLY: Place the appropriate number that describes the variate abstracted or five suriety in the board on a minimum of 100 plants. Comparative data should be determined from varieties entered in the same trial. Royal Horticultural Society or any recognized color standard may be used to determine plant colors, designate system used: Please answer all questions for your variety; lack of response may delay progress of your application.  1. KIND:  1			WIIDAT (Trucum	i shb.)		•
ADDRESS (Street and No. or RD No., Cip., Sunt, and Exp Code)  PARTY MANGE  VARIETY MANGE  We in dy  TEMPORARY On EXTENDIORY ADDRESS (CHATTON)  Place a zero in the first box (e.g. [1] [1] or [1] ]) when number is either 99 or less or 9 or less respectively. Data for quantitative plant character should be based on a minimum of 100 plants. Comparative data abould be determined from varieties entered in the same trial. Royal Horizoultural Society or any recognized color standard many be used to determine plant colors; designate system used:  Pleaze answer all questions for your variety; lock of response may delay progress of your application.  1. KIND:  1	NAME OF APPLICANT(S)	· · · · · · · · · · · · · · · · · · ·			FOR OFFICIAL USE ONLY	
VARIETY NAME Wendy  TEMPORARY OR EXPRENDENT LIPESONATION  Place a zero in the first box (e.g. [3] [3] or [3] [2]) when number is either 99 or less or 9 or less respectively. Data for quantities plant character should be based on a minimum of 100 plants. Comparative data should be determined from varieties entered in the same trial. Royal Horticultural Society or any recognized color standard may be used to determine plant oring; designate system used:  Please answer all questions for your variety; lack of response may delay progress of your application.  1. KIND:  1			· · · · · · · · · · · · · · · · · · ·			
PLEASE READ ALL INSTRUCTIONS CAREFULLY: Place the appropriate number that describes the varietal character of this variety in the boxes below. Place a zero in the first box (e.g. [3-3] or [a-1]) when number is either 99 or less or 9 or less respectively. Data for quantitative plant characters should be based on a minimum of 100 plants. Comparative data should be determined from warteless entered in the same trial. Royal Horticultural Society or any recognized color standard may be used to determine plant colors; designate system used:  1. KIND:  1	ADDRESS (Street and No. or RD No.,	City, State, and Zip Code)			200501	102
PLEASE READ ALL INSTRUCTIONS CAREFULLY: Place the appropriate number that describes the varietal character of this variety in the boxes below. Place a zero in the first box (e.g. [2 2 3] or [3 3]) when number is either 99 or less or 9 or less respectively. Data for quantitative plant characters should be based on annimum of 100 plants. Comparative data should be determined from varieties entered in the same trial. Royal Horticultural Society or any recognized color standard may be used to determine plant colors; designate system used:  Please answer all questions for your variety; lack of response may delay progress of your application.  1. KIND:  1					VARIETY NAME Wendy	
Place a zero in the first box (e.g. [3 [3 ] ar [3 ]) when number is either 99 or less or 9 or less respectively. Data for quantitative plant characters should be bettermined from varieties entered in the same trial. Royal Horticultural Society or any recognized color standard may be used to determine plant colors; designate system used: Please answer all questions for your variety; lack of response may delay progress of your application.  1. KIND:  1			·	•	TEMPORARY OR EXPERIMENTAL SD 9 7 W 6 (	designation ) 4
1 1=Common 2=Durum 3=Club 4=Other (SPECIFY):  2 VERNALIZATION:  2 1=Spring 2=Winter 3=Other (SPECIFY):  3 COLEOPTILE ANTHOCYANIN:  1=Absent 2=Present  4. JUVENILE PLANT GROWTH:  2 1=Prostrate 2=Semi-erect 3=Erect  5. PLANT COLOR (boot stage):  2 1 = Yellow-Green 2 = Green 3 = Blue-Green  6. FLAG LEAF (boot stage):  2 1 = Erect 2 = Recurved 1 = Not Twisted 2 = Twisted  7. EAR EMERGENCE:  Harding	Place a zero in the first box (e a minimum of 100 plants. Co may be used to determine plan	.g. 0 9 9 or 0 9 ) when omparative data should be determent colors; designate system used:	number is either 99 or less or 9 or nined from varieties entered in th	or less respectively. he same trial. Royal I	Data for quantitative plant charac	ters should be based on
2. VERNALIZATION:  2	1. KIND:					
2	1	1=Common	2=Durum	3=Club	4=Other (S)	PECIFY):
3. COLEOPTILE ANTHOCYANIN:  1=Absent 2=Present  4. JUVENILE PLANT GROWTH:  2 1=Prostrate 2=Semi-erect 3=Erect  5. PLANT COLOR (boot stage):  2 1 = Yellow-Green 2 = Green 3 = Blue-Green  6. FLAG LEAF (boot stage):  2 1 = Erect 2 = Recurved 1 = Not Twisted 2 = Twisted  7. EAR EMERGENCE:  Number of Days Earlier Than  Harding	2. VERNALIZATION	N:	, <del>**</del>		· · · · · · · · · · · · · · · · · · ·	·
1=Absent 2=Present  4. JUVENILE PLANT GROWTH:  2    1=Prostrate 2=Semi-erect 3=Erect  5. PLANT COLOR (boot stage):  2    1 = Yellow-Green 2 = Green 3 = Blue-Green  6. FLAG LEAF (boot stage):  2    1 = Erect 2 = Recurved 1 = Not Twisted 2 = Twisted  7. EAR EMERGENCE:    Number of Days Earlier Than   Harding   Hardin	2	1=Spring	2=Winter	3=Other (SI	PECIFY):	
4. JUVENILE PLANT GROWTH:  2 1=Prostrate 2=Semi-erect 3=Erect  5. PLANT COLOR (boot stage):  2 1 = Yellow-Green 2 = Green 3 = Blue-Green  6. FLAG LEAF (boot stage):  2 1 = Erect 2 = Recurved 1 = Not Twisted 2 = Twisted  7. EAR EMERGENCE:  Number of Days Earlier Than  Harding	3. COLEOPTILE AN	THOCYANIN:	· · · · · · · · · · · · · · · · · · ·			
2 1=Prostrate 2=Semi-erect 3=Erect  5. PLANT COLOR (boot stage):  2 1 = Yellow-Green 2 = Green 3 = Blue-Green  6. FLAG LEAF (boot stage):  2 1 = Erect 2 = Recurved 1 = Not Twisted 2 = Twisted  7. EAR EMERGENCE:  Number of Days Earlier Than  Harding		1=Absent	2=Present			
5. PLANT COLOR (boot stage):  2	4. JUVENILE PLANT	GROWTH:		· · · · · · · · · · · · · · · · · · ·		
1 = Yellow-Green 2 = Green 3 = Blue-Green  6. FLAG LEAF (boot stage):  2	2	1=Prostrate	2=Semi-erect	3=Erect		
6. FLAG LEAF (boot stage):  2	5. PLANT COLOR (I	boot stage):				· · · · · · · · · · · · · · · · · · ·
2	2	1 = Yellow-Green	2 = Green	3 = Blue-Gre	een	·
7. EAR EMERGENCE:  Harding  Number of Days Earlier Than  *	6. FLAG LEAF (boot	stage):		···········		
Number of Days Earlier Than*	2	I = Erect	2 = Recurved		1 = Not Twisted	2 = Twisted
Number of Days Earlier Than*	7. EAR EMERGENC	E:				
LILL-54		Number of Days Earlie	Harding er Than			
0 0 Number of Days Later Than Expedition	0 0	Number of Days Later	ThanExpedit	ion		*

8. ANTHER COLOR:	
1 = Yellow 2 = Purple	
9. PLANT HEIGHT (from soil to top of head, excluding awns):	
cm Taller Thannone	•
0 8 cm Shorter Than Expedition	•
	Relative to a PVPO-Approved Commercial Variety Grown in the Same Trial
10. STEM:	
A. ANTHOCYANIN	D. INTERNODE (SPECIFY NUMBER)
1 1= Absent 2=Present	1 = Hollow 2=Semi-solid , 3=Solid
B. WAXY BLOOM	E. PEDUNCLE
1=Absent 2=Present	1=Absent 2=Present
C. HAIRINESS (last internode of rachis)	cm Length
1=Absent 2=Present	
11. HEAD (at Maturity):	
A. DENSITY	C. CURVATURE
2=Middense 3= Dense	2 1 = Erect 2 = Inclined 3 = Recurved
B. SHAPE	D. AWNEDNESS
1 = Tapering 2= Strap 3 = Clavate 4 = Other (SPECIFY):	1 = Awnless 2 = Apically Awnletted 3 = Awnletted 4 = Awned
12. GLUMES (at Maturity):	
A. COLOR	C. BEAK
1 = White  2 = Tan	3 1 = Obtuse 2 = Acute
3 = Other (SPECIFY) :	3 =Acuminate
B. SHOULDER	D. LENGTH
1 = Wanting 2 = Oblique 3 = Rounded 4 = Square 5 = Elevated 6 = Apiculate	1 = Short 2 = Medium (ca. 7mm) (ca. 8mm) 3 = Long (ca. 9mm)

12.	GLUMES	(at Maturity) Co	ntinued:					
	E. W	IDTH					200500	102
	2	1 = Narrow (ca. 3 = Wide (ca. 4)		Medium (ca. 3.5m)	m)			
13.	SEED:		<u></u> .				· · · · · · · · · · · · · · · · · · ·	····
	A. SH	IAPE			C. BR	USH		
	3	1 = Ovate	2 = Oval	3 = Elliptical	1	1=Short	2=Medium	3=Long
					1	1 = Not Collared	2 = Collared	•
	B. CF	IEEK			D. CR	EASE		
	1	1=Rounded	2=Angu <u>l</u> ar		1	1 = Width 60% or less 2 = Width 80% or less 3 = Width Nearly as W	of Kernel	••• •••
					1	<ul> <li>1 = Depth 20% or less</li> <li>2 = Depth 35% or less</li> <li>3 = Depth 50% or less</li> </ul>	of Kernel	
	E. Col	or	-		G. PH	ENOL REACTION (se	e instructions):	
	1	1=White 4= OTHER (Sp	2= Amber ecify)	3= Red	3	1 = Ivory 3 = Light Brown 5 = Black	2 = Fawn 4 = Dark Brow	n
	F. TE	XTURE				•		
	1	1=Hard	2=Soft					
14.	DISEASE:	(0=Not Tes	ted; 1=Susce	ptible; 2=Resis	tant; 3	B=Intermediate; 4=To	lerant)	· · · · · · · · · · · · · · · · · · ·
		. I	PLEASE INDIC	ATE THE SPEC	IFIC RAC	CE OR STRAIN TESTE	D	
	2	Stem Rust (Puo	ccinia graminis (	L sp. <i>tritici)</i>	3	Leaf Rust (Puccinia re	condita f. sp. tritic	<b>D</b>
	1	Stripe Rust (Pu	iccinia striiform	is)		Loose Smut (Ustilago	tritici)	
	2	Tan Spot (Pyre	nophora tritici-r	repentis)	0	Flag Smut (Urocystis a	igropyri)	
	0	Halo Spot <i>(Sele</i>	enophoma donac	cis)	0	Common Bunt (Tilleti	a tritici or T. laevis	5)
	0	Septoria nodoru	m (Glume Blot	ch)	0	Dwarf Bunt (Tilletia c	ontroversa)	· ·
	0	Septoria avenae	(Speckled Lea	f Disease)	0	Karnal Bunt (Tilletia	indica)	
	0	Septoria tritici (	Speckled Leaf	Blotch)	0	Powdery Mildew (Ery	siphe graminis f. s	p. <i>tritici)</i>
	1	Scab (Fusariun	n spp.)		0	"Snow Molds"		•
								1

				200500 100
14.	Diseas	e (Continued) (0=Not Tested; 1=Suscepti	ble; 2=	Resistant; 3=Intermediate; 4=Tolerant)
		PLEASE INDICATE TH	HE SPECI	FIC RACE OR STRAIN TESTED
	0	"Black Point" (Kernel Smudge)	0	Common Root Rot (Fusarium, Cochliobolus and Bipolaris spp.)
	0	Barley Yellow Dwarf Virus (BYDV)	0	Rhizoctonia Root Rot (Rhizoctonia solani)
	1	Soilborne Mosaic Virus (SBMV)	0	Black Chaff (Xanthomonas campestris pv. translucens)
	O	Wheat Yellow (Spindle Streak) Mosaic Virus	0	Bacterial Leaf Blight (Pseudomonas syringae pv. syringae)
	2	Wheat Streak Mosaic Virus (WSMV)		Other (SPECIFY)
<del>reger</del> su*		Other (SPECIFY)		Other (SPECIFY)
		Other (SPECIFY)		Other (SPECIFY)
		Other (SPECIFY)		Other (SPECIFY)
15.	INSECT:	(0=Not Tested; 1=Susceptible; 2=Resis	stant; 3	=Intermediate; 4=Tolerant)
		PLEASE SPECIFY B	IOTYPE	(where needed)
	1	Hessian Fly (Mayetiola destructor)		Other (SPECIFY)
		Stem Sawfly (Cephus spp.)		Other (SPECIFY)
	0	Cereal Leaf Beetle (Oulema melanopa)		Other (SPECIFY)
	0	Russian Aphid (Diuraphis noxia		Other (SPECIFY)
<b>.</b> k	0	Greenbug (Schizaphis graminum)		Other (SPECIFY)
·	0	Aphids		Other (SPECIFY)

16. ADDITIONAL INFORMATION ON ANY ITEM ABOVE, OR GENERAL COMMENTS

# Exhibit D Wendy (SD97W604) Additional Description of the Variety

The following additional descriptive information is presented:

- 1) Release notice of Wendy.
- 2) Table 2: Yield, test weight, heading, and plant height data of Wendy in the South Dakota Crops Performance Testing (CPT) variety trial in 23 locations-years.
- 3) Table 3: Yield and test weight performance data of Wendy in the South Dakota Crops Performance Testing (CPT) variety trial in 2004.
- 4) Table 4: Yield and test weight performance data of Wendy in the South Dakota Crops Performance Testing (CPT) variety trial in 2003.
- 5) Table 5: Yield and test weight performance data of Wendy in the South Dakota Crops Performance Testing (CPT) variety trial in 2002.
- 6) Table 6: Milling and baking data of Wendy in the South Dakota Crops Performance Testing.
- 7) Table 7: Quality parameters of selected lines grown in South Dakota environments and tested at the Hard Winter Wheat Quality Laboratory, USDA/ARS, Manhattan, KS.

To:301 504 5291

### South Dakota Agricultural Experiment Station South Dakota State University Plant Science Department

### Release of 'Wendy' Hard White Winter Wheat

'Wendy' hard white winter wheat (Triticum aestivum L.) was developed by the South Dakota Agricultural Experiment Station and released to seed producers in August 2004. Wendy was released on the basis of its white grain color, good noodle quality, excellent winter survival, earliness, and high yield potential in South Dakota and the northern Great Plains region. Wendy has been named in memory of Wendy Wickersham, secretary for South Dakota Wheat Inc. from 2003-2004.

Wendy was derived from the cross \$D89333/'Abilene' made in 1992. SD89333 is an unreleased experimental line from South Dakota State University with the pedigree 'Gent'/'Siouxland'. Wendy was developed by means of the bulk breeding method. The cross (coded X92259) was advanced to the F3 generation as a bulk. Seed harvested from the F3 bulk was sorted for white kernel color in 1995. The bulk of selected white kernels was coded X92259W and was grown in the greenhouse in 1996. Single heads were harvested from the selected F<sub>4</sub> bulk and planted in the field as head-rows in fall 1996. Wendy was derived as an F<sub>4,5</sub> line selected by S.D. Haley in 1997. Wendy was evaluated as SD97W604 in the South Dakota Early Yield Trial nursery in 1998. It was advanced beyond the Preliminary Yield Trial to the South Dakota Advanced Yield Trial in 1999 due to superior performance. It was tested in the South Dakota Crop Performance Testing (CPT) Variety Trial between 2000 and 2004, in the Northern Regional Performance Nursery during 2001 and 2002, and in the Southern Regional Performance Nursery in 2004. Breeder seed of Wendy originated from a composite of 200 F<sub>10-11</sub> head-rows selected in 2002 based on visual uniformity and white kernel color purity.

Wendy is an awned, white-glumed, early maturing, semi-dwarf hard white winter wheat. Wendy has green foliage at anthesis. The spike is tapered, inclined, and mid-dense. The glume size is medium, and the glume shoulder has a wanting shape. The beak is medium in length with an acuminate tip. Kernels are white, hard textured, and elliptical in shape with a collarless short brush, rounded cheeks, and a shallow crease.

Wendy is early maturing (146 d to heading from 1 Jan.), 1 d earlier than 'Expedition' (PI 629060), 3 d earlier than 'Wesley' (PI605742), and 6 d earlier than 'Harding' (PI608049) in South Dakota yield trials. Plant height (75 cm) of Wendy was similar to Wesley and 15 cm less than Harding. The winter survival of Wendy is good to excellent (similar to Harding). It has a short coleoptile (60 mm; 86% of Wesley; 75% of Expedition; and 67% of Harding) and excellent straw strength (similar to Wesley). Wendy has fair to good resistance to pre-harvest sprouting (3.7 score; 1 = highly resistant to 9 = highly susceptible), similar to Trego (P1612576) (3.4), higher than Millennium (PI613009) (5.0) and Jerry (PI632433) (8.0), and lower than Expedition (1.9) and Nekota (PI584997) (1.0). Because of its level of sprouting resistance, Wendy will be best adapted to areas west of the Missouri river in South Dakota,

In 31 site-years of testing in the South Dakota CPT, yield of Wendy (3534 kg ha<sup>-1</sup>) was higher than 'Harding' (3468 kg ha<sup>-1</sup>), 'Wesley' (3451 kg ha<sup>-1</sup>), 'Expedition' (3413 kg ha<sup>-1</sup>), 'Arapahoe' (PI518591) (3395 kg ha<sup>-1</sup>), and 'Trego' (3405 kg ha<sup>-1</sup>), and lower than 'Millennium' (3625 kg ha<sup>-1</sup>) and 'Jagalene' (3663 kg ha<sup>-1</sup>). Wendy had similar volume weight to Trego (764 kg m<sup>-3</sup>) higher than Wesley (745 kg m<sup>-3</sup>) and lower than Jagalene (775 kg m<sup>-3</sup>).

Wendy has moderate resistance to stem rust (caused by *Puccinia graminis* f. sp. trittci.) and has been postulated to carry Sr24 and Sr31 based on tests conducted by the USDA Cereal Disease Laboratory, St. Paul, MN. Wendy has been confirmed to be homogeneous for the IBL.1RS wheat-rye translocation based on SDS-PAGE gel analysis. It is moderately susceptible to leaf rust (caused by *Puccinia triticina* Eriks.) and is tolerant to tan spot (caused by *Pyrenophora tritici-repentis* (Died.) Drechs.). Wendy is moderately susceptible to wheat streak mosaic virus, and is susceptible to the Great Plains biotype of Hessian fly [Mayetiola destructor (Say)]. Wendy has exhibited intermediate reaction to wheat soil-borne mosaic virus.

Composite milling and bread baking properties of Wendy were determined in 2001 and 2002 in cooperative baking tests conducted by the USDA/ARS Hard Winter Wheat Quality Laboratory in Manhattan, KS. Milling scores were very good while baking scores were poor. Relative to the hard white check cultivars Trego and 'Nuplains' (PI605741), Wendy had medium-sized kernels (28.2 versus 29.0 and 25.7 mg, respectively). Flour extraction of Wendy, Trego, and Nuplains was 672, 667 and 673 g kg<sup>-1</sup>, respectively. Flour ash of Wendy was similar to Trego (4.2 g kg<sup>-1</sup>) and slightly lower than Nuplains (4.3 g kg<sup>-1</sup>). Flour protein of Wendy (112 g kg') was lower than Nuplains (115 g kg') and higher than Trego (110 g kg'). In bread baking tests, bake absorption of Wendy (585 g kg<sup>-1</sup>) was lower than both Nuplains (615 g kg<sup>-1</sup>) and Trego (612 g kg<sup>-1</sup>), while its loaf volume was similar to Nuplains (0.88 L) and higher than Trego (0.85 L). Wendy had lower mixograph tolerance than both Trego and Nuplains (1.0 versus 2.5 and 3 scores, respectively; 0 = unacceptable to 6 = excellent scale). Wendy had lower mixograph mix time (3.7 min) than Trego (4.1 min) and Nuplains (5.1 min). Wendy was tested in the Wheat Quality Council in 2003. It was slightly better than Crimson (PI601818), and had good milling and acceptable bread baking quality characteristics. Wendy has low grain polyphenol oxidase (PPO) levels. In tests conducted by the Asian Products Collaborative (APC) Project, coordinated by the U.S. Wheat Associates and the Wheat Marketing Center, in 2003, Wendy was similar or better than Australian standards for Korean steamed buns, modified Chinese northern type steamed bread, and Chinese raw noodles.

Breeder seed of Wendy originated from a composite of 200  $F_{10.11}$  head-rows selected in 2002 based on visual uniformity and white kernel color purity. Wendy has been uniform for all morphological characters (such as maturity and plant height) during the last four generations of increase. Wendy contains 0.14% hard red grain. It also contains red and tall white off-types in the frequency of 0.05% and 0.01%, respectively.

The South Dakota Foundation Seed Stocks Division (Plant Science Department, South Dakota State University, Brookings, SD) has foundation seed of Wendy available to seed producers for planting during fall 2004. Seed classes will be Breeder, Foundation, Registered, and Certified. Wendy will be submitted for plant variety protection under P.L. 910577 with the certification option.

<u>Approval</u>

Director, South Dakota Agricultural Experiment Station

Date

Table 2. Agronomic data for Wendy (2001 - 2003)

Entry	_ID	GY	TW	SS	HD	НТ
1	ALLIANCE	48.5	57.6	7.0	157	30.9
2	ARAPAHOE	50.1	58.2	7.0	159	33.0
3	CRIMSON	46.7	59.4	7.2	161	35.1
4	EXPEDITION	48.9	58.3	7.2	156	30.9
5	FALCON	51.0	57.1	7.0	160	29.5
6	HARDING	48.3	58.5	7.2	161	35.2
7	JERRY	49.4	57.7	6.8	162	35.2 35.3
8	MILLENNIUM	51.6	59.5	6.8	160	33.6
9	NEKOTA	48.1	58.7	7.0	157	29.8
10	NUPLAINS	46.3	59.2	6.1	161	29.6
11	RANSOM	46.6	57.0	7.1	162	29.6 35.5
12	SD92107-3	48.1	58.1	7.0	161	
13	SD92107-5	49.2	58.9	7.0 7.1	162	34.1
14	SD97049	47.5	57.8	6.5	158	34.7
15	SD97250	47.8	58.2	7.2	159	29.5
16	WENDY	48.7	58.7	6.2		32.7
17	SD97W609	49.3	58.7	6.4	156	27.5
18	TANDEM	48.8	59.7		157	28.3
19	TREGO	48.0	59.7 59.0	7.2	159	34.1
20	WAHOO	49.6		6.5	158	29.5
21	WESLEY		57.1 59.0	6.6	159	31.4
Mean	WEGLET	49.7	58.2	7.1	157	28.0
CV%		48.7	58.4	6.9	159	31.8
LSD 0.05		13.2	3.2	19.4	0.7	4.8
L3D 0.03		1.6	0.5	0.5	0.5	0.6

Table 3. Grain Yield and Test Weight data for the 2004 Crops Performance Testing (CPT) Variety Trial

	7000			1		Grain Y	Grain Yield (bu/a)						i i
Entry	2004		(	1	D. Lakes						Mater		I wy (ID/bu)
Lated y	Average	e z-yr	3-yr	Brookings	J	Highmore	Platte	Selby	Sturgis	Wall	water- town	Winner	Average
SD97538	65	61		103	55	28	ç	ţ					200
SD92107-5	63	59	53	96	52	2 - 2	3 6	· -	82	53	47	58	29
SD00W024	63		<b>}</b>	8 6	7 6 2	9 7	0 1	1.7	27	61	42	49	59
SD97059-2	62	61		8 8	3 5	- 6	<u>.</u>	20	28	59	44	53	26
SD97394-1	63	5		0 6		82	62	72	24	52	52	52	α C
SD98102	7 G	08		55 G	58	7.7	62	75	29	47	46	12	, 10 10 10 10 10 10 10 10 10 10 10 10 10
70100	70	8	55	35	54	78	69	73	30	49	53	- 67	20.00
WAHOO	62	<u>c</u>	7	700	ç	í	·				}	2	9
HARDING	5	- 04	t c	200	53	8/	63	20	26	57	37	52	22
TERRY	5 6	9 6	70	93 0	46	9/	72	89	27	56	45	52	ο α γ
COOC	- i	28	51	106	46	79	62	71	29	53	<u>د</u> ج	4 6	9 6
SD00258	61			103	47	73	99	. 2	5 6	3 6		40	28
MILLENNIUM	09	9	54	100	47	7.4	80	2 6	+ 0	ر ا	4/	25	58
SD99073	90			5	7.	1.4	8 8	2 8	R7 ;	47	46	55	59.
					5	2	70	99	24	49	46	51	29
JAGALENE	29	90	55	83	ğ	42	į	í	,				
NE99533-4	59	;	8	3 6	9 4	 10	22	7.	24	52	39	09	09
5097250	0 4	ų u	,	Q	22	5	09	29	29	20	44	20	9
AL TANCE	3 4	8 [	ີດີ	82	54	74	63	64	27	46	56	49	50
יייייייייייייייייייייייייייייייייייייי	20 1	26	51	83	53	75	64	71	29	46	50	2 7	9 6
SD00111	58			84	55	92	59	7	? ?	7	9 4		6G
SD97380-2	58	28		06	7.	1	8 4	- 1	<b>5.</b>	40	46	48	59
		1		}	5		S.C.	/9	24	45	48	47	59
SD00032	22			G	53	74	41	Ç.	;				
TANDEM	57	ያሂ	04	8	3 8	- i	/6	68	19	46	53	49	59
SD97W671-1	2 6	3 1	2	4 6	53	Z,	63	65	26	51	42	46	90
WECLEY	5 6	5 [	į	<b>X</b>	55	92	53	29	21	47	41	46	20
יירטקני	00	/6	51	96	47	72	09	58	25	48	42	20	0 0
SUS/WEUS	99	28	52	98	20	29	58	69	22	5. 4	38	60 4	90
- KEGO	22	56	51	82	49	68	59	63	27	. 9	8 8	† <del>1</del>	8 8
								}	į	3	<b>,</b>	<i>/</i> c	.0
NEKOLA	52	55	20	98	51		58	09	28	45	24	7	
SDUUWU41	54			78	45	73	56	65	2	70	- K	, t	n c
ARAPAHOE	54	56	51	6/	45		57	74	i 6	2 5	2.3	g !	
CRIMSON	54	55	48	08	73		. 1	- 0	17	40	34	4/	58
EXPEDITION	53	, r.	. r	6 6	3 4		70	00	77	48	41	38	
SD97W604	53	7 9	- 6	- 6	÷.		28	62	21	47	36	40	
	3	õ	S.	00	74		48	92	25	45	39	20	59
MEAN	59	28	52	91	51		61	89	26	49	44	. 0	
LSD .05	3.5	2.1	1.8	16.4	13.1		10.4	6.9	7.2	5.7	43.5		
% CA%	12.2	11.6	13.4	12.9	18.2	7.2	12.1	7.2	17.4	. 6		73	7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7
/										- 5	5		o <u>.</u>

Table 4. Yield and test weight performance data of Wendy (SD97W604) in the South Dakota CPT in 2004

	GY	TW			<del>-</del> -			rain Y	ield b	u/ac				··
<u>ID</u>	AVG	AVG	Bis	Oel	Mar	Hay		DLP			Win	Pla	Hig	Ken
SD97W604	60.8	60.9	57.2	66.1	68.0								51.2	
SD97W609	60.2	60.3	51.8	68.7	64.3		44.3		_			-	60.8	
JAGALENE	60.1	60.9	55.2	76.6		61.5		-	90.2	_			57.4	89.2
MILLENNIUM	59.7	60.7	55.6	62.6	68.6	62.5						68.9	56.8	77.0
WAHOO	59.3	58.2	53.5	70.7	75.2	53.8	44.9	36.6			48.7	65.4	56.8	80.6
SD97538	58.3	59.3	49.4	68.9	72.0	55.5	47.4	36.0				64.7	50.9	
WESLEY	58.2	59.2	51.9	64.9	70.8	61.5	40.7	36.4	83.1	44.3	43.1	66.1		83.6
SD98102	58.1	59.6	49.3	67.9	67.8	47.1	46.5	36.3	84.0	44.3	50.0	71.3	55.4 54.1	79.9
EXPEDITION	58.0	59.8	54.5	72.3	68.0	58.4	43.8	37.5	78.6	39.5	46.9	63.5	54.1 51.2	79.2
SD97059-2	58.0	58.9	52.7	61.4	65.8	44.3	44.1	36.3	98.9	43.0	46.9	62.7		82.0
SD97380-2	57.9	58.9	49.9	68.5	67.1	57.3	46.1	39.4	89.4	37.0	46.6	59.9	53.6	86.4
SD99W015	57.4	59.5	51.8	66.5	65.3	54.1	46.0	36.5	76.9	41.3	49.3		56.0	78.0
FALCON	57.3	59.4	56.5	66.8	61.5	50.7	43.5	35.9	83.3	41.1	49.3	70.8	54.4	76.1
ARAPAHOE	57.2	59.3	52.8	64.5	67.8	57.2	45.8	38.0	85.3	35.9	47.3	60.3 59.6	55.7 57.0	85.1
SD97049	56.6	58.8	51.3	66.6	65.9	50.4	42.7	31.4	87.4	43.4	43.5	69.4	57.0	77.9
SD97W671-1	56.2	60.2	52.8	71.0	63.6	43.9	43.7	35.6	77.9	43.4	52.8		56.4	70.7
SD97088	56.2	59.8	48.1	65.0	66.6	48.2	44.4	33.8	88.5	45.4 45.4	52.6 43.8	62.9	54.6	73.0
TREGO	56.0	60.2	56.9	69.8	69.6	55.0	43.2	33.5	75.6	38.3	43.6 42.9	65.1	53.6	72.2
ALLIANCE	55.4	58.5	52.6	70.7	69.3	54.0	43.6	37.4	70.9	43.6	41.3	61.3	51.9	73.5
SD92107-5	55.3	60.3	49.7	66.3	66.8	47.3	42.2	31.4	83.3	44.4	43.5	55.5 59.4	53.4	72.8
HARDING	54.8	59.9	52.8	68.0	64.6	45.9	39.5	37.6	88.4	39.2	43.5	55.3	54.8	74.3
NEKOTA	54.7	59.9	50.5	69.6	60.0	51.6	41.8	34.4	79.0	41.7	48.5		54.5	70.3
SD97250	54.4	59.2	50.6	66.8	61.4	51.3	44.3	34.5	79.3	36.9	46.5 45.9	61.7	49.3	68.5
JERRY	54.3	59.7	50.4	56.6	59.0	47.3	40.0	36.2	86.5	41.4	45.9 46.2	62.5	53.0	66.6
CRIMSON	53.8	61.2	53.0	68.7	60.0	38.2	40.2	42.5	84.8	41.4	45.8	57.0 55.0	56.9	74.8
TANDEM	53.5	61.2	52.4	65.6	67.5	48.9	41.5	35.4	74.7	42.4 42.9		55.2	48.4	66.3
SD92107-3	53.4	59.5	48.0	60.7	65.0	43.8	43.4	35.6	86.7	42.9 37.6	45.4	51.0	52.2	65.1
NUPLAINS	53.2	61.2	48.6	65.0	58.5	49.7	41.3	37.9			41.8	56.0	53.2	69.4
AP502CL		57.5	50.7	72.8	60.6	60.3	41.3 37.6	36.3	86.6 63.3	43.9	44.9	51.2	50.4	60.1
RANSOM		58.7	47.0	54.5	55.5	44.9	39.2	33.5		39.5	42.6	61.1	40.2	71.2
Mean		59.7	51.9	66.8	65.4	52.2	43.1	36.0	77.6	36.8	45.0	49.0	48.4	63.8
CV%		90.1	8.0	6.5	4,3	10.9			83.2	40.9	46.0	61.9	53.4	75.4
-SD (0.05)			5.8	6.1	4.3 3.9		5.9	11.6	8.8	12.1	8.5	10.7	8.1	9.8
-0- (0.00)			J.0	0.1	ა.ყ	8.0	3.6	6.0	10.3	6.9	5.9	9.4	6.2	2.0

Table 5. Grain yield and test weight means for 2002 South Dakota winter wheat crop performance testing (CPT) variety trial

		33		6	irc	l'a-	1~~	, Tan	, I -		- I -	·	. 1 =			,	,		-						_																
MEAN	Ē	25		1		ı	ì					1	- 1		f	1	1	1					- 1	27		1										l					
JEAN M	<u>ર</u>	46	45	43	42	42	42	42	45	1.1	4	41	- 4	4	9	40	40	40	40	9	3	8 6	Ş	5 6	39	99	က္က	38	38	38	38	37	37	36	36	36	36	35	4		-
BRON	TW	29	90	8	22	28	8	8	28	۲. ا	32.	3	9	20	6	9	28	22	09	8	28	28	20	8 09	28	09	8	29	28	29	27	27	20	8	22	61	09	8	29	က	65
380		- 1	i		1	1	1				-			1	1	1			1	1	i i			5, 5		ı			1	l	1			1			Į.				ŀ
J A L	0000	- 1	ı				1	1	1			1		1	1		1		1	ı	f	1		57			ì		1	ĺ											f
PLA .	33.6	- [					1	1	1			1	1			Ī	1	ł				1	1			1	1	1			' '								1		•
WAT			- 1	1	- 1						1		1	ļ	1		[	i	i	1	ł		1	55	i				1			1	- 1	ı							
WAT W				_ ]				i	ŀ	1		1		ŀ	li							1		1		]				- 1	- 1		- 1		- 1	ı		- 1			
WAL W	1000	- 1	- 1			. !				Į	1												ļ	56 4		i I		- 1		ſ	ı	- 1		- i		- 1		.		- 1	
0.000			_	- 1	- 1	ł	ľ		1						- 1		- 1	ŀ		ŀ				1 1	ı	- 1		ı		- 1	- 1		- 1		- 1	- 1	- 1				
N WAL	¥	$\perp$		╛													ľ		ļ					. 1	1			ļ	ĺ		- 1	- 1			- 1	ſ	-		ļ	- [	
NIM Z	ě			-		-	- 1	- 1	ł				f	- 1	- 1			ŀ	- 1	-		- 1	i				- 1	ı			-	- 1		1				- 1			
NIM .	á	1				- !		- 1	Ì	- !				ı		ı	- 1		ſ	- 1	ı			27		- 1		- !	- 1			ļ		ı		-	ŀ		1		
SEL	Ä		-	1	ı		- 1	- }	ı			ĺ	- 1			ı	Ţ	- 1		ŀ		- 1		99	- 1	ľ						1	1			- 1			- 1		- 1
SEL	28	35	3 6	2 5	7 2	9	27	32	28	29	83	3	26	30	25	77	23	177	3	77	17	78	29	25	87	26	3 5	7	47 6	2 8	7 6	4 6	; 2	- 6	3 8	2 8	3 8	7 5	17	- (	16
HIG. TW	09	90	200	2 2	3 2	n c	8 8	25 5	56	28	22	8	20	28	<u>ت</u> ا و	8	3 8	2 2	200	20 00	3	6	29	23	3	6	3 8	3 5	0 0	18	3	3 6	8	8 8	8 8	3 8	0 8	3 8	ñ,	=	=
) E &	33	32	27	1 %	200	2 2	3 8	3	55	3	8	67	3	2 3	4, 6	9 6	3 8	3 2	\$ 8	₹ 8	8	77	3	হ্য হ	77	3 8	77	3 5	* o	200	3 8	3 2	5 8	27	27 6	77	2 2	S S	0 0	1 c	=
	59	62	61	57	α̈́	3 8	3 6	8	1 6	2	20 2	٥	3 8	5 6	2 2	3 6	3 2	o d	5 6	8 8	8 8	3 8	20	8	8 8	8	3 8	9 6	3 2	3 6	5 6	2 5	3 6	3 4	3 2	3 2	3 6	3 2	3 0	1 0	7
SY.	46	22	47	43	47	17	7	‡ =	‡ 5	ئ ئ	7 5	200	7 0	t c	3 r	2 4	2 6	5 4	2 6	2 10	,	7 0	04	4 6	2 2	1 5	3 5	3 5	7 7	000	3 8	3 8	8	7	2	3 5	2 5	3 5	2 ~	- 0	ō
		25	22	53	53	27	, L	5 6	2 5	2 2	4 6	1 8	à la	5 6	5 6	2 4	2 2	5 2	ן א	3 6	5 4	8 3	4 5	2 2	7 6	o a	3 12	3 12	27	7 7	. 29	2 42	2 2	52	1 6	70	2 2	3 15	3 0	1 0	7
ر و۲			1	i	1		ļ	- 1	1	Į.		-	ſ	1								ı		4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4				47			2 4						44		4	7	-
	i		<u> </u>	l	1	1	1	1			0 8	3 8	3 &	- «	3 6	65	1 6	3 8	1 8	3 8	3 2	2 2	5 6	2 8	3 8	3 8	1 6	1 6					-	1	83	2 5	3 6	1 6	1 0	1 6	1
	ſ		ı	1		1	i	1	-	- 1		- 1	1	-			1	-	1	ĺ		- 1		- 1			1	-	1	ı	1		1						16		ł
	- 1	- 1				1				1		- 1			1							1	- 1	8 6		1		- 1	1	1	1	1	F		1						ı
	- !	- 1	ì							1	-						1						1	3 8		1			ł	1	ı	1			1	1			1	0	,
	4		(7)		(1)	(7)	(*)	(6)		,	1 (**	) (*		l c	(6)	E.		2	0	7	1	10	1 (	) (r.	0	1 6	2	6	2	e	100	7	m	2	22	2	2	1 (4)		-	
	17406	ا					Σ			ш	7604)					ER										甲							z								
i i	COAN IUM/406	JACALENE	7W604	FALCON	WESLEY	SD97457	MILLENNIUM	TANDEM	WAHOO	ARAPAHOF	Ch (SD97W604	OTA	SD97250	90	STANTON	RONT	ALLIANCE	SD92107-3	SD92107-5	NW98S059	7049	2137	HARDING	SD97W650	NUPLAINS	<b>AVALANCHE</b>	3102	3306	7088	ځ	3ER	7432	NUHORIZON	RANSOM	SCOUT66	CRIMSON	SD98W198	_	.05		
2		S 6	SOS	FAL	WE	SD9	MILL	TAN	WAF	ARA	ਠ	NEK	SD9	TRE	STA	N H	ALLI	SD9,	SD9.	NWS	SD9	213	HAR	SD9	NU	AVA	SD98102	SD96306	SD97088	JERI	JAGGER	SD97432	N	RAN	SCO	양	SD9	Mean	LSD .05	% S	
<u>;</u>	=   _	<b>1</b>	3	2	၈	23	13	ω	4	2	8	2	22	32	9	8	_	19	20	37	77	42	8	35	28	29	26	24	25	မွ	16	27	ਲ	7	17	15	36				

Table 6. Milling and baking data of Wendy in the South Dakota Crops Performance Testing

ID	Nuplains	Wendy	Trego	Wesley
Grain volume weight, kg hl-1	46.4	46.3	46.2	45.0
Percent large kernels, %	46.8	63.3	65.0	76.1
Kernel wieght, mg	25.7	28.2	29.0	32.1
SKCS kernel hardness	72.3	73.2	70.2	63.6
Flour ash, g kg-1	4.3	4.2	4.2	4.5
Flour extraction, g kg-1	673	672	667	688
Flour protein content, g kg-1	115	112	110	121
Water Absorbtion, g kg-1	615	585	612	631
Make mix time, min	5.1	3.7	4.1	7.8
Mixograph tolerance	3.0	1.0	2.5	4.5
Loaf volume, L	0.88	0.88	0.85	0.86
Crumb grain score	4.6	3.5	4.0	4.9

Table 7. Quality parameters of selected lines grown in south Dakota environments and tested at the Hard Winter Wheat Quality Laboratory, USDA/ARS, Manhattan, KS.

	i	Single	Kernel		Flour Ch	aracter	-		Mixing		Baki	ng .
	ID	SK_H	SK_W	F_YLD	F_ASH	F_PRO	F_ABS	M_ABS	M_MT	M_TOL	L_VOL	B_MT
	Arapahoe	61.5	28.8	68.4	0.40	10.7	60.9	60.8	3.9	4.0	805	5.7
	Crimson	82.8	27.3	66.8	0.42	10.5	61.2	61.1	2.6	3.0	905	3.4
	Expedition	57.8	32.8	71.1	0.34	10. <del>9</del>	61.0	61.1	5.5	5.0	795	7.4
2	Harding	76.8	29.5	66.1	0.38	10.9	61.1	61.1	3.9	4.0	910	5.6
0	Nuplains	74.3	26.6	69.3	0.40	10.4	61.1	61.3	2.6	3.0	825	4.5
0	Trego	66.9	30.7	69.0	0.38	10.2	61.2	60.9	3.3	3.0	825	3.7
1	WENDY	69.8	28.5	67.9	0.40	10.5	57.1	60.6	2.6	1.0	890	3.4
	Nursery St. Dev.	7.4	2.8	1.4	0.04	0.4	1.3	0.9	8.0	1.0	45	1.1
	Nursery C.V. %	11.6	9.1	2.0	8.84	4.1	2.1	1.4	22.3	28.3	5	23.3
Ш	Nursery Mean	64.3	30.2	68.5	0.40	10.7	60.5	60.8	3.8	3.5	834	4.8
	Subset Mean	70.0	29.2	68.4	0.39	10.6	60.5	61.0	3.5	3.3	851	4.8
	Arapahoe	69.8	23.4	63.3	0.56	12.8	62.0	64.3	4.1	3.0	925	5.3
l I	Crimson	83.3	20.0	57.4	0.55	12.9	63.1	63.9	3.1	4.0	950	4.3
	Expedition	61.2	27.0	66.3	0.47	12.2	63.0	63.2	5.5	5.0	900	9.0
2	Harding	67.0	22.2	62.7	0.53	13.2	63.4	64.5	4.5	4.0	990	5.8
0	Nuplains	70.4	22.7	65.3	0.47	12.7	61.8	63.6	3.4	3.0	935	4.8
0	Trego	73.4	24.4	64.4	0.46	11.8	61.2	62.7	3.4	2.0	875	3.5
2	WENDY	76.7	22.0	66.6	0.44	11.8	59.8	59.6	2.8	1.0	875	3.2
	Nursery St. Dev.	6.8	1.3	2.5	0.04	0.5	1.0	1.2	0.6	1.0	50	1.3
	Nursery C.V. %	9.4	5.8	3.9	7.29	4.1	1.7	1.9	15.1	28.8	- 5	23.6
Ш	Nursery Mean	71.8	23.1	63.7	0.51	12.4	62.0	62.9	4.1	3.6	927	5.4
	Subset Mean	71.7	23.1	63.7	0.50	12.5	62.0	63.1	3.8	3.1	921	5.1
	Arapahoe	71.3	29.7	66.3	0.31	12.3	63.6	63.4	4.0	4.0	845	5.0
ll	Crimson	77.8	27.2	65.1	0.31	12.8	62.2	64.3	2.1	3.0	915	3.1
2	Expedition	66.4	29.6	67.6	0.30	12.7	63.6	64.1	5.8	6.0	925	8.3
0	Harding	79.2	29.3	65.2	0.34	12.8	63.1	64.3	4.1	4.0	970	4.6
0	WENDY	64.2	28.0	65.8	0.31	12.7	60.7	60.1	3.5	1.0	905	3.9
	Nursery St. Dev.	4.9	1.7	1.6	0.02	0.5	0.9	1.3	0.7	1.0	40	1.2
	Nursery C.V. %	6.9	5.7	2.4	5.45	3.9	1.5	2.0	17.2	23.8	4	21.4
_	Nursery Mean	70.2	29.1	66.8	0.32	12.4	62.8	63.6	4.3	4.2	906	5.6
	Subset Mean	44.3	18.9	40.5	1.28	8.5	38.0	38.4	6.3	7.2	569	7.9
2	Arapahoe	67.5	27.3	66.0	0.42	11.9	62.2	62.8	4.0	3.7	858	5.3
0	Crimson	81.3	24.9	63,1	0.43	12.1	62.2	63.1	2.6	3,3	923	3.6
1	Expedition	61.8	29.8	68.4	0.37	11.9	62.5	62.8	5.6	5.3	873	8.2
2	, Harding	74.3	27.0	64.7	0.42	12.3	62.5	63.3	4.2	4.0	957	5.3
0	WENDY	70.2	26.2	66.8	0.38	11.7	59.2	60.1	3.0	1.0	890	3.5
	Nursery Mean	68.8	27.5	66.3	0.41	11.9	61.8	62.4	4.0	3.8	889	5.3
	Subset Mean	71.0	27.0	65.8	0.40	12.0	61.7	62.4	3.9	3.5	900	5.2

SK\_H = Single kernel hardness (scale of 1-100); SK\_W = Single kernel weight (mg); F\_YLD = Flour yield (percent of total product); F\_ASH = Flour ash (percent at 14 % moisture basis); F\_PRO = Flour protein (percent at 14 % moisture basis); F\_ABS = Flour absorption (14 % moisture basis); M\_ABS = Mixograph absorption (14 % moisture basis); M\_MT = Mixograph peak mix time (minutes); M\_TOL = Mixograph tolerance; L\_VOL = Loaf volume; B\_MT = Baking mix time (minutes).

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE	The following statements are made in accordance with the Privacy Act of 1974 (5 U.S.C. 552a) and the Paperwork Reduction Act (PRA) of 1995.	
EXHIBIT E STATEMENT OF THE BASIS OF OWNERSHIP	Application is required in order to determine if a plant variety protection certificate is to be issued (7 U.S.C. 2421). Information is held confidential until certificate is issued (7 U.S.C. 2426).	
1. NAME OF APPLICANT(S)	2. TEMPORARY DESIGNATION OR EXPERIMENTAL NUMBER	3. VARIETY NAME
South Dakota Agricultural Experiment Station	SD97W604	Wendy
4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP, and Country)	5. TELEPHONE (include area code)	6. FAX (include area code)
South Dakota State University Ag Hall 129	(605) 688-4149	(605 688-6065
Brookings SD 57007	7. PVPO NUMBER	<u> </u>
	2	005 0 0 102
Is the applicant (individual or company) a U.S. national or U.S. based     if no, give name of country	company?	YES NO
10. Is the applicant the original owner?	NO If no, please answer one of the	following:
a. If original rights to variety were owned by individual(s), is (are) the	original owner(s) a U.S. national(s)?	
- ·	NO If no, give name of country	
b. If original rights to variety were owned by a company(ies), is(are) th	ne original owner(s) a U.S. based compan	y?
☐ YES ☐	NO If no, give name of country	
11. Additional explanation on ownership (if needed, use reverse for extra	space):	
		·
	<u>.</u>	

#### PLEASE NOTE:

Plant variety protection can be afforded only to owners (not licensees) who meet one of the following criteria:

- 1. If the rights to the variety are owned by the original breeder, that person must be a U.S. national, national of a UPOV member country, or national of a country which affords similar protection to nationals of the U.S. for the same genus and species.
- 2. If the rights to the variety are owned by the company which employed the original breeder(s), the company must be U.S. based, owned by nationals of a UPOV member country, or owned by nationals of a country which affords similar protection to nationals of the U.S. for the same genus and species.
- 3. If the applicant is an owner who is not the original owner, both the original owner and the applicant must meet one of the above criteria.

The original breeder/owner may be the individual or company who directed final breeding. See Section 41(a)(2) of the Plant Variety Protection Act for definition.

According to the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to compete this information collection is estimated to average 10 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, and maritat or familial status. (Not all prohibited bases apply to all programs). Persons with disabilities who require alternative means for communication of program information (braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint, write the Secretary of Agriculture, U.S. Department of Agriculture, Washington, D.C. 20250, or call 1-800-245-6340 (voice) or (202) 720-1127 (TDD). USDA is an equal employment opportunity employer.